

GUIDELINES FOR VAGINAL BIRTH AFTER PREVIOUS CAESAREAN BIRTH

This guideline has been prepared and reviewed by the Clinical Practice Obstetrics Committee and approved by Executive and Council of the Society of Obstetricians and Gynaecologists of Canada.

PRINCIPAL AUTHORS

Marie-Jocelyne Martel, MD, FRCSC, Saskatoon SK
Catherine Jane MacKinnon, MD, FRCSC, Brantford ON

CLINICAL PRACTICE OBSTETRICS COMMITTEE

Catherine Jane MacKinnon, MD, FRCSC, Brantford ON
Marc-Yvon Arseneault, MD, FRCSC, Montreal QC
Elias Bartellas, MD, FRCSC, St. John's NL
Yvonne M. Cargill, MD, FRCSC, Ottawa ON
Sue Daniels, RN, Dartmouth NS
Tom Gleason, MD, FRCSC, Edmonton AB
Stuart Iglesias, MD, Gibsons BC
Michael C. Klein, MD, CCFP, Vancouver BC
Marie-Jocelyne Martel, MD, FRCSC, Saskatoon SK
Ann Roggensack, MD, Kingston ON
Ann Kathleen Wilson, BHSc, RM, Ilderton ON

SPECIAL CONTRIBUTOR

Gregory A. Davies, MD, FRCSC, Kingston ON

Abstract

Objective: To provide evidence-based guidelines for the provision of a trial of labour (TOL) after Caesarean section.

Outcome: Fetal and maternal morbidity and mortality associated with vaginal birth after Caesarean (VBAC) and repeat Caesarean section.

Evidence: MEDLINE database was searched for articles published from January 1995 to February 2004, using the key words "vaginal birth after Caesarean (Cesarean) section." The quality of evidence is described using the Evaluation of Evidence criteria outlined in the Report of the Canadian Task Force on the Periodic Health Exam.

Recommendations:

1. Provided there are no contraindications, a woman with 1 previous transverse low-segment Caesarean section should be offered a trial of labour after Caesarean with appropriate discussion of maternal and perinatal risks and benefits. The process of informed consent with appropriate documentation should be an important part of the birth plan in a woman

with a previous Caesarean section. (II-2B)

2. The intention of a woman undergoing a TOL after Caesarean should be clearly stated and documentation of the previous uterine scar should be clearly marked on the prenatal record. (II-2B)
3. For a safe labour after Caesarean section, the woman should deliver in a hospital where an immediate Caesarean section is available. The woman and her health-care provider must be aware of the hospital resources and the availability of obstetric, anaesthesia, pediatric, and operating-room staff. (II-2A)
4. Each hospital should have a written policy in place regarding the notification and/or consultation for the physicians responsible for a possible immediate Caesarean. (III B)
5. Continuous electronic fetal monitoring of women attempting a TOL after Caesarean is recommended. (II-2A)
6. Suspected uterine rupture requires urgent attention and expedited laparotomy in order to attempt to decrease maternal and perinatal morbidity and mortality. (II-2A)
7. Oxytocin augmentation is not contraindicated in women undergoing a TOL after Caesarean. (II-2A)
8. Medical induction of labour with oxytocin may be associated with an increased risk of uterine rupture and should be used carefully after appropriate counselling. (II-2B)

Key Words

Vaginal birth after Caesarean, trial of labour, uterine rupture, induced labour, oxytocin, prostaglandins, misoprostol

These guidelines reflect emerging clinical and scientific advances as of the date issued and are subject to change. The information should not be construed as dictating an exclusive course of treatment or procedure to be followed. Local institutions can dictate amendments to these opinions. They should be well documented if modified at the local level. None of the contents may be reproduced in any form without prior written permission of SOGC.

FOR INFORMATION ON THE SELF-DIRECTED LEARNING EXERCISE, SEE PAGE 684.

9. Medical induction of labour with prostaglandin E₂ (dinoprostone) is associated with an increased risk of uterine rupture and should not be used except in rare circumstances after appropriate counselling. (II-2B)
10. Prostaglandin E₁ (misoprostol) is associated with a high risk of uterine rupture and should not be used as part of a TOL after Caesarean. (II-2A)
11. A foley catheter may be used safely to ripen the cervix in a woman planning a TOL after Caesarean. (II-2A)
12. The available data suggest that a trial of labour in women with more than one previous Caesarean section is likely to be successful but is associated with a higher risk of uterine rupture. (II-2B)
13. Multiple gestation is not a contraindication to a TOL after Caesarean. (II-2B)
14. Diabetes mellitus is not a contraindication to TOL after Caesarean. (II-2B)
15. Suspected fetal macrosomia is not a contraindication to a TOL after Caesarean. (II-2B)
16. Women delivering within 18 to 24 months of a Caesarean section should be counselled about an increased risk of uterine rupture in labour. (II-2B)
17. Postdatism is not a contraindication to a TOL after Caesarean. (II-2B)
18. Every effort should be made to obtain the previous Caesarean section operative report to determine the type of uterine incision used. In situations where the scar is unknown, information concerning the circumstances of the previous delivery is helpful in determining the likelihood of a low transverse incision. If the likelihood of a lower transverse incision is high, TOL after Caesarean can be offered. (II-2B)

Validation: These guidelines were approved by the Clinical Practice Obstetrics and Executive Committees of the Society of Obstetricians and Gynaecologists of Canada.

J Obstet Gynaecol Can 2004;26(7):660–70.

BACKGROUND

This document will review the contraindications to and maternal and fetal risks of trial of labour (TOL) after Caesarean birth and make recommendations for achieving vaginal birth after Caesarean (VBAC) safely. Delivery by Caesarean section occurs in 15% to 25% of births.¹⁻⁵ In 2000 and 2001, the Caesarean section rate in Canada was 21.2%.⁶ The most frequent indications for Caesarean section are previous Caesarean section, dystocia, malpresentation, and non-reassuring fetal status.^{7,8} In any given region, the rate of birth by Caesarean section and the rate of VBAC tend to be inversely related.⁴ Schell first reported VBAC in 1923, describing the successful vaginal delivery of 34 infants in 23 mothers with previous Caesarean deliveries.⁹

A trial of labour after Caesarean should be considered in women who present for prenatal care with a history of previous Caesarean birth.¹⁰⁻¹² In certain situations, TOL after Caesarean will be contraindicated³ and a repeat Caesarean section will be advised, but in most cases, successful vaginal birth can be achieved safely for both mother and infant.¹³⁻¹⁵ Women and

their health-care providers will need to discuss the risks and benefits of VBAC when planning the birth.

A Canadian consensus statement on VBAC was published in 1985 and Clinical Practice Guidelines were published by the Society of Obstetricians and Gynaecologists of Canada (SOGC) in 1997.³ This document updates the 1997 SOGC Guidelines, with articles published from January 1995 to February 2004, obtained by searching the MEDLINE database using the key words “vaginal birth after Caesarean (Cesarean) section.” The data is limited by 3 important factors: first, there are no randomized trials of TOL versus elective repeat Caesarean section (ERCS); second, adverse maternal or perinatal outcomes are rare and large study populations are necessary to observe a significant difference in maternal and perinatal outcomes; and, finally, the woman’s choice to attempt a TOL after Caesarean is heavily influenced by her health-care provider and local resources, often leading to selection bias in published reports.^{12,16}

The level of evidence and quality of the recommendations in this guideline have been determined using the criteria described by the Canadian Task Force on the Periodic Health Examination (Table).¹⁷

TRIAL OF LABOUR VERSUS ELECTIVE REPEAT CAESAREAN SECTION

The success rate of trial of labour after Caesarean ranges between 50% and 85%.^{3,4,14,18-21} In a study examining 1776 women undergoing TOL after Caesarean, the overall success rate was 74%.¹⁴ A Canadian study reported similar results quoting a success rate of 76.6%.² Predictors of successful VBAC include non-recurring indication for Caesarean birth, such as malpresentation (odds ratio [OR], 1.9; 95% confidence interval [CI], 1.0–3.7)²² or gestational hypertension (OR, 2.3; 95% CI, 1.0–5.8),²² and a previous vaginal delivery (OR, 1.8; 95% CI, 1.1–3.1),²² where success rates are as high as 82%.^{1,22,23} When the previous Caesarean birth was for dystocia, failure to progress, or cephalopelvic disproportion, some studies found the rates of successful VBAC comparable,^{24,25} while others reported or lower-than-expected rates.^{14,18,22,26}

In 1996, McMahon *et al.* published a report of maternal morbidity in TOL compared to ERCS in Nova Scotia from 1986 to 1992.¹ In an examination of 3249 women undergoing a TOL and 2889 women who delivered by ERCS, the risk of major complications (hysterectomy, uterine rupture, and operative injury) was almost doubled in the TOL group (1.6% vs. 0.8%: OR, 1.8; 95% CI, 1.1–3.0).¹ Complications like puerperal fever, transfusion and abdominal wound infection were comparable. When comparing those who had a successful TOL to those who required a repeat Caesarean section after failed TOL, the risks were greater of operative injury (3.0% vs. 0.1%: OR, 5.1; 95% CI, 2.5–10.7) and fever (8.0% vs. 3.5%: OR,

1.5; 95% CI, 1.3–1.8) in the failed TOL group.¹ Hibbard *et al.* also reported a greater rate of complication in those who had attempted a TOL and failed.²⁷

In 1999, Ragoth *et al.* reviewed 17 613 TOL and 11 433 ERCS deliveries.²⁰ The rates of hysterectomy (relative risk [RR], 0.36; 95% CI, 0.23–0.56), febrile morbidity (RR, 0.65; 95% CI, 0.55–0.77), and thromboembolic complications (RR, 0.52; 95% CI, 0.34–0.78) were less in the TOL group than in the ERCS group.²⁰ There is less blood loss with a successful VBAC (OR, 0.50; 95% CI, 0.3–0.9)²⁷ and a shorter hospital stay with a more rapid recovery and return to full activity.

Rosen *et al.* also reported that the risk of febrile morbidity is lower in women who attempt a TOL after Caesarean (OR, 0.5; 95% CI, 0.5–0.6) and is lowest in those who succeed (OR, 0.2; 95% CI, 0.2–0.2), compared to ERCS, but is increased in those who attempt a TOL and ultimately deliver by Caesarean section (OR, 2.0; 95% CI, 1.7–2.5).²⁸

An examination of 16 938 Finnish women who had undergone a Caesarean delivery found that previous Caesarean section is associated with an increased risk of ectopic pregnancy (RR, 1.28), placenta previa (RR, 3.89), and abruptio placenta (RR, 2.41).²⁹ A repeat Caesarean section has been associated with an increase in the risk of placenta previa (OR, 1.59; 95% CI, 1.21–2.08)³⁰ and placenta accreta in subsequent pregnancies.³¹

A meta-analysis published in 2000 demonstrated that the overall risk of perinatal death is increased in those attempting a TOL (OR, 1.71; 95% CI, 1.28–2.28).³² The risks of perinatal mortality and severe morbidity are directly related to uterine rupture as a sentinel event. If uterine rupture occurs, the risk of perinatal mortality and severe morbidity are increased (see below). The risk of suspected neonatal sepsis is greater in those attempting a TOL but appears to be confined to the group of TOL who fail and require a repeat Caesarean section (OR, 4.8; 95% CI, 2.6–9.0).³³ If an ERCS is chosen, the risk of respiratory problems in the newborn is increased compared to those

who have a successful VBAC (6% vs. 3%: OR, 2.3; 95% CI, 1.4–3.8).³³

CONTRAINDICATIONS TO VAGINAL BIRTH AFTER CAESAREAN SECTION

The following situations are contraindications to a TOL after Caesarean:

1. previous classical or inverted “T” uterine scar^{3,13};
2. previous hysterotomy or myomectomy entering the uterine cavity^{3,19};
3. previous uterine rupture^{3,19};
4. the presence of a contraindication to labour, such as placenta previa, malpresentation³;
5. the woman declines a TOL after Caesarean and requests ERCS.^{3,19}

RECOMMENDATION

1. **Provided there are no contraindications, a woman with 1 previous transverse low-segment Caesarean section should be offered a trial of labour after Caesarean with appropriate discussion of maternal and perinatal risks and benefits. The process of informed consent with appropriate documentation should be an important part of the birth plan in a woman with a previous Caesarean section. (II-2B)**

PLANNING A TRIAL OF LABOUR AFTER CAESAREAN SECTION

The woman and her health-care provider must decide together whether an appropriate situation exists for considering a TOL after Caesarean. The evaluation and discussion should address the issues outlined below and these should be well documented in the prenatal record or chart.

Quality of Evidence Assessment	Classification of Recommendations
<p>The quality of evidence reported in these guidelines has been described using the Evaluation of Evidence criteria outlined in the Report of the Canadian Task Force on the Periodic Health Exam.</p> <p>I: Evidence obtained from at least one properly randomized controlled trial.</p> <p>II-1: Evidence from well-designed controlled trials without randomization.</p> <p>II-2: Evidence from well-designed cohort (prospective or retrospective) or case-control studies, preferably from more than one center or research group.</p> <p>II-3: Evidence obtained from comparisons between times or places with or without intervention. Dramatic results in uncontrolled experiments (such as the results of treatment with penicillin in the 1940s) could also be included in this category.</p> <p>III: Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.</p>	<p>Recommendations included in these guidelines have been adapted from the ranking method described in the Classification of Recommendations found in the Report of the Canadian Task Force on the Periodic Health Exam.</p> <p>A. There is good evidence to support the recommendation that the condition be specifically considered in a periodic health examination.</p> <p>B. There is fair evidence to support the recommendation that the condition be specifically considered in a periodic health examination.</p> <p>C. There is poor evidence regarding the inclusion or exclusion of the condition in a periodic health examination, but recommendations may be made on other grounds.</p> <p>D. There is fair evidence to support the recommendation that the condition not be considered in a periodic health examination.</p> <p>E. There is good evidence to support the recommendation that the condition be excluded from consideration in a periodic health examination.</p>

DOCUMENTATION OF PREVIOUS UTERINE INCISION

Documentation of the location and type of uterine incision used during the previous Caesarean section is ideal.³ In most cases, this information can be obtained by reviewing the operative record from the previous surgery. Other information in this record, such as the indication for the Caesarean section and the opinion of the previous surgeon, may be helpful in counselling as well. The fact that the record has been reviewed and that no contraindications to a TOL after Caesarean are present should be documented clearly on the prenatal record.³⁴ If the operative record is not available, the scar is considered “unknown.” Review of the operative report from previous hysterotomy or myomectomy should be documented in detail.

RECOMMENDATION

2. **The intention of a woman undergoing a TOL after Caesarean should be clearly stated and documentation of the previous uterine scar should be clearly marked on the prenatal record. (II-2B)**

FACILITIES AND RESOURCES

A trial of labour after Caesarean is always associated with a risk of uterine rupture, however small. For this reason, a TOL after Caesarean should only be considered in a hospital where provisions for performing an immediate Caesarean section are available.^{13,34,35} Facilities providing VBAC should have a policy in place to manage such parturients so that all resources are mobilized promptly if an intrapartum emergency occurs.²³ The availability and time required for obstetric, anaesthesia, and pediatric services to attend such an emergency should be fully discussed with the woman. Women who live in areas where local hospitals cannot offer immediate Caesarean section should be offered the opportunity for transfer to a facility where this service is available, in order to permit a TOL after Caesarean.¹³ The members of the team who could be called urgently in the case of an intrapartum complication (anaesthesia, pediatric, and obstetric services) should be notified that the woman is in hospital and in labour and their availability confirmed.

A woman with a previous Caesarean section should be advised against planned out-of-hospital labour and birth.

RECOMMENDATION

3. **For a safe labour after Caesarean section, the woman should deliver in a hospital where an immediate Caesarean section is available. The woman and her health-care provider must be aware of the hospital resources and the availability of obstetric, anaesthesia, pediatric, and operating-room staff. (II-2A)**
4. **Each hospital should have a written policy in place regarding the notification and/or consultation for the physicians responsible for a possible immediate Caesarean. (III B)**

MATERNAL MONITORING

The woman planning a TOL after Caesarean should have appropriate monitoring in labour. The presence of a devoted birth attendant is ideal. Progress of labour should be assessed frequently as there is some evidence that prolonged or desultory labour is associated with an increased risk of failure and uterine rupture.^{19,36,37} Epidural analgesia is not contraindicated.^{7,19,34,38}

FETAL MONITORING

Continuous electronic fetal monitoring in labour is recommended for all women attempting a TOL after Caesarean.^{19,34,39} The most reliable first sign of uterine rupture is a non-reassuring fetal heart tracing.³⁴ This may be sudden in onset and may not be related to contractions.⁴⁰

RECOMMENDATION

5. **Continuous electronic fetal monitoring of women attempting a TOL after Caesarean is recommended. (II-2A)**

POSTPARTUM EVALUATION

Routine digital exploration of the Caesarean section scar postpartum is not necessary, except when signs or symptoms suggest uterine rupture.⁴¹

UTERINE RUPTURE

Uterine rupture, the most serious complication of a TOL after Caesarean, is defined as complete separation of the myometrium with or without extrusion of the fetal parts into the maternal peritoneal cavity requiring emergency Caesarean delivery or postpartum laparotomy.^{19,42} It is an uncommon complication of VBAC, but is associated with significant maternal and perinatal morbidity and mortality.^{1,7} The most common sign or symptom of uterine rupture is non-reassuring fetal heart rate monitoring.^{18,20,43} Other clinical signs include the cessation of contractions, loss of the presenting part on vaginal examination, abdominal pain, vaginal bleeding, hematuria, or maternal cardiovascular instability.^{16,44}

The type and location of the previous uterine incision helps to determine the risk of uterine rupture. The incidence of uterine rupture is 0.2% to 1.5% in a woman who attempts labour after a transverse lower-uterine-segment incision^{14,16,18,27,45} and 1% to 1.6% after a vertical incision in the lower uterine segment.⁴⁶⁻⁴⁹ The risk is 4% to 9% with a classical or “T” incision; and for this reason, a TOL after Caesarean is contraindicated in these situations.^{16,19,30} Shimonovitz *et al.* found the risk of uterine rupture after 0, 1, 2, and 3 VBAC deliveries to be 1.6%, 0.3%, 0.2%, and 0.35%, respectively, indicating that the risk of uterine rupture decreases after the first successful VBAC.⁵⁰

Since uterine rupture is a rare event, a realistic appraisal of potential maternal and perinatal risks is difficult to accomplish

outside of large series, literature reviews, or meta-analyses. The most important published reports in this area are discussed below as well as those applicable to the Canadian population.

In 1991, Rosen *et al.* performed a meta-analysis including 10 studies examining 4617 women who had a TOL after Caesarean compared to 3831 women who had ERCS births.²⁸ The rate of uterine rupture was similar in the two groups: TOL (0.18%), ERCS (0.19%, $p=0.5$). There was no difference in the rate of maternal death (0.028% vs. 0.024%) or perinatal death (0.3% vs. 0.4%) in the TOL compared to the ERCS group.²⁸

In 2000, Mozurkewich and Hutton published a meta-analysis of 15 studies examining 28 813 women undergoing a TOL compared to ERCS between 1989 and 1999.³² There was an increased rate of uterine rupture (0.39% vs. 0.16%; OR, 2.1; 95% CI, 1.45–3.05) and perinatal mortality (0.58% vs. 0.28%; OR, 1.71; 95% CI, 1.28–2.28) in the TOL group. The rates of maternal mortality and low 5-minute Apgar scores were not different.³²

In 2002, Keiser *et al.* reviewed the incidence and consequences of uterine rupture in Nova Scotia from 1988 to 1997.⁵¹ Among 4516 women undergoing a TOL, 18 (0.39%) uterine ruptures were documented over 10 years. All women underwent laparotomy and there were no maternal deaths. Of those who had a uterine rupture, 3 women underwent hysterectomy, 10 required transfusion, and 5 suffered a cystotomy. After excluding lethal anomalies, there was 1 perinatal death (0.02%) and 6 neonates with severe asphyxia (0.13%).⁵¹

In 2002, Bujold *et al.* examined the risk factors for serious neonatal morbidity associated with 23 cases of uterine rupture among 2233 women attempting a TOL (rate 1.03%).⁵² Nine neonates (0.4%) had a $pH < 7.0$ (severe metabolic acidosis), 3 (0.13%) were diagnosed with hypoxic ischemic encephalopathy, and 1 (0.04%) died.⁵² The presence of placental or fetal part extrusion at laparotomy was associated with severe metabolic acidosis ($p < 0.001$).⁵² Other variables (e.g., induction, birth weight, use of epidural) did not demonstrate an association with uterine rupture. Even in situations where very rapid decision to delivery times were recorded, some cases of perinatal acidosis could not be avoided.⁵²

Smith *et al.* published a large series of 15 515 women undergoing a TOL after Caesarean compared to 9014 who underwent ERCS between 1992 and 1997.⁵³ The rate of perinatal death in the TOL group was 0.129%, 11.6 times higher than that of the ERCS group (OR, 11.6; 95% CI, 1.6–86.7).⁵³ Smith compared this to the risk of perinatal death in other common obstetrical situations: TOL compared to multiparous woman in labour (OR, 2.2; 95% CI, 1.3–3.5) and TOL compared to nulliparous woman in labour (OR, 1.3; 95% CI, 0.8–21).⁵³

In 2003, Chauhan *et al.* published a review of data on the maternal and perinatal complications of uterine rupture in those attempting a TOL after Caesarean.⁵⁴ Examining 142 075 trials

of labour revealed an overall rate of uterine rupture of 0.62%.⁵⁴ The rate of maternal death was 0.002%; hysterectomy, 0.09%; transfusion, 0.18%; and genitourinary tract injury, 0.08%.⁵⁴ In this study, the rate of neonatal acidosis was 0.15%, and the rate of perinatal death was 0.04%.⁵⁴ Oxytocin was involved in 43% of the uterine ruptures in this series.⁵⁴

The data indicate that the relative risk of uterine rupture, maternal morbidity, and perinatal mortality or severe morbidity is increased in those undergoing a TOL after Caesarean compared to ERCS, but that the absolute risk remains very low.

The treatment of suspected uterine rupture is immediate laparotomy after maternal stabilization and anaesthesia. Urgent intervention is mandatory to obtain the best possible outcome for both mother and fetus. Once the fetus is delivered, maternal hemorrhage must be arrested, and if the uterus cannot be salvaged, hysterectomy may be required.

Although the risk of uterine rupture has been found to be increased in situations of prolonged labour with augmentation,⁵⁵ when Phelan *et al.* retrospectively examined the patterns of uterine activity before uterine rupture, no association with frequency or intensity of contractions could be discerned.⁴⁰

In 1996, Rozenberg *et al.* examined ultrasonographic measurement of the lower uterine segment's myometrial thickness at 36 to 38 weeks' gestation as a predictor of uterine rupture and found that if the lower segment thickness was less than 3.5 mm, the risk of uterine rupture or dehiscence was 11.8%; and if the measurement was greater than 3.5 mm, the risk of uterine rupture was minimal.⁵⁶ However, the incidence of uterine rupture in this population was 2.3%, significantly greater than the usually quoted 1%, and therefore the positive predictive value of this test in clinical practice will be much lower.⁵⁶ In a follow-up open study, Rozenberg *et al.* found that the use of the lower-uterine-segment measurement helped clinicians select women for a TOL after Caesarean.⁵⁷ The rate of successful VBAC for those with 1 previous Caesarean section did not change, but was increased in those with 2 previous Caesarean sections.⁵⁷ These findings will need to be confirmed in further randomized studies before ultrasonography can be used to make a decision about the safety of a TOL after Caesarean.

RECOMMENDATION

- Suspected uterine rupture requires urgent attention and expedited laparotomy in order to attempt to decrease maternal and perinatal morbidity and mortality. (II-2A)**

OXYTOCICS AND TRIAL OF LABOUR AFTER CAESAREAN SECTION

AUGMENTATION

In 1987, Flamm *et al.* performed a multicentre examination of 485 women who received oxytocin to augment their spontaneous labour in a planned TOL after Caesarean.⁵⁸ No increase in the

risk of uterine rupture, maternal morbidity, or perinatal morbidity or mortality was detected.⁵⁸ Zelop *et al.* supported the same conclusion about the risk of uterine rupture with augmentation in a 1999 study (OR, 2.3; 95% CI, 0.8–7.0).⁵⁹ Goetzl *et al.* examined the relationship between the dose of oxytocin used and the risk of uterine rupture in women undergoing a TOL after Caesarean.⁶⁰ No significant association was detected between exposure to oxytocin and the risk of uterine rupture.⁶⁰ Careful surveillance of the woman for progress of labour is required, especially when the diagnosis of dystocia is being considered.^{19,34} There are insufficient studies examining the use of other agents to augment labour, such as prostaglandins, and their safety in a TOL after Caesarean.

INDUCTION

In 2000, Ravasia *et al.* reviewed the risk of uterine rupture in women undergoing an induction TOL after Caesarean.⁶¹ In 575 women with a previous Caesarean section, labour was induced with prostaglandin E₂ gel (n=172), intracervical foley catheter (n=129), or amniotomy and/or oxytocin (n=274).⁶¹ Outcomes were compared to those women undergoing a TOL with spontaneous labour. The risk of uterine rupture was not increased in those who underwent either amniotomy/oxytocin or foley catheter induction, but was significantly increased in those who underwent a prostaglandin E₂ induction (p=0.004).⁶¹

Also in 2000, Sanchez-Ramos *et al.* performed a meta-analysis looking at the efficacy and safety of prostaglandin E₂ for cervical ripening in women with a previous Caesarean section and found it to be effective and not associated with an increased risk of uterine rupture (OR, 1.46; 95% CI, 0.96–2.22) compared to spontaneous labour.⁶²

In 2003, Delaney and Young reported the examination of 3746 women with a prior Caesarean delivery who underwent either induced or spontaneous labour.⁶³ They found that induced labour was associated with a greater risk of early postpartum hemorrhage (7.3% vs. 5.0%: OR, 1.66; 95% CI, 1.18–2.32), Caesarean delivery (37.5% vs. 24.2%: OR, 1.84; 95% CI, 1.51–2.25), and admission to a neonatal intensive care unit (13.3% vs. 9.4%: OR, 1.69; 95% CI, 1.25–2.29).⁶³ There was a trend towards a higher rate of uterine rupture, but this was not statistically significant (0.7% vs. 0.3%, p=0.128).⁶³

In another retrospective study of 560 women, the rate of uterine rupture in women whose labour was induced with oxytocin was 2%, with prostaglandin was 2.9%, and with both was 4.5%.⁵⁹

Up to 2001, there was conflicting data on the risk of induction of labour with prostaglandin E₂. Several other smaller studies reported that it appeared to be safe, effective, and not associated with an increased risk of uterine rupture.^{45,64-66}

In the largest study published to date, conducted by Lydon-Rochelle *et al.*, the incidence of uterine rupture was reviewed retrospectively in 20 095 women with a previous Caesarean

section and reported as follows: elective repeat Caesarean section (no labour) 0.16%; spontaneous labour 0.52% (RR, 3.3; 95% CI, 1.8–6.0); labour induced without prostaglandin 0.77% (RR, 4.9; 95% CI, 2.4–9.7); and labour induced with prostaglandin 2.45% (RR, 15.6; 95% CI, 8.1–30.0).⁶⁷

As for all inductions, the indication for induction in a woman undergoing a planned TOL after Caesarean should be compelling and documented. The possibility that the use of oxytocin and/or prostaglandin for induction of labour in women considering a TOL after Caesarean may be associated with an increased risk of uterine rupture and its sequelae must be discussed with the parturient. The absolute risks of uterine rupture are low, but the relative risks (especially with the use of prostaglandin E₂ compared to spontaneous labour) are greater.

MISOPROSTOL

Misoprostol has been proposed as an effective and economical agent for cervical ripening and induction.⁶⁸ In 1998, Sciscione *et al.* reported a case of uterine rupture in a woman with 2 previous Caesarean sections after misoprostol was administered as a cervical ripening agent.⁶⁹ A number of small series reported a risk of uterine rupture from 0% to 11.7% with misoprostol in women undergoing a TOL after Caesarean.^{43,70-73} Blanchette *et al.* compared prostaglandin E₂ to misoprostol in women undergoing induction TOL after Caesarean and found them to be equally effective, but misoprostol was associated with a higher incidence of uterine rupture (18.8% compared to no ruptures in the prostaglandin E₂ group).⁷⁴ The numbers in all of these studies are small and it is difficult to draw meaningful conclusions. Until further randomized studies are completed, misoprostol should be discouraged as a method of induction or cervical ripening in women with previous Caesarean delivery.^{74,75}

CERVICAL PREPARATION

In situations where delivery is indicated and the cervix is unfavourable, a TOL after Caesarean can be considered. Various methods of cervical ripening have been examined. In a cohort study published in 2002, Ben-Aroya *et al.* compared women undergoing a trial of labour after Caesarean section in 3 situations: spontaneous labour (n=1432), prostaglandin cervical ripening (n=55), and cervical ripening by foley catheter (n=161).⁷⁶ There was a significantly higher rate of dystocia (30.4% vs. 11.6%, p<0.01) and repeat Caesarean section in the second stage (49.1% vs. 35.2%, p<0.01) in the foley catheter group compared to the control group.⁷⁶ There was no difference in the rate of uterine rupture, fetal distress, or Apgar scores.⁷⁶ In a Canadian study published in 2004, Bujold *et al.* compared the rate of uterine rupture in 1807 women who presented in spontaneous labour, 417 induced with amniotomy

with or without oxytocin, and 255 induced with transcervical foley catheter.⁷⁷ The rate of successful vaginal birth was 78% in the spontaneous group, 77.9% in the amniotomy group, and 55.7% in the transcervical foley group ($p < 0.001$).⁷⁷ However, the rates of uterine rupture did not differ significantly: 1.1%, 1.2%, and 1.6%, respectively ($p = 0.81$).⁷⁷ These data support the use of the foley catheter for cervical ripening of an unfavourable cervix in women undergoing a TOL after Caesarean.

RECOMMENDATIONS

7. **Oxytocin augmentation is not contraindicated in women undergoing a TOL after Caesarean. (II-2A)**
8. **Medical induction of labour with oxytocin may be associated with an increased risk of uterine rupture and should be used carefully after appropriate counselling. (II-2B)**
9. **Medical induction of labour with prostaglandin E₂ (dinoprostone) is associated with an increased risk of uterine rupture and should not be used except in rare circumstances after appropriate counselling. (II-2B)**
10. **Prostaglandin E₁ (misoprostol) is associated with a high risk of uterine rupture and should not be used as part of a TOL after Caesarean. (II-2A)**
11. **A foley catheter may be used safely to ripen the cervix in a woman planning a TOL after Caesarean. (II-2A)**

SPECIAL CIRCUMSTANCES

MORE THAN 1 PREVIOUS LOW TRANSVERSE CAESAREAN SECTION

Several authors have assessed the rate of successful VBAC and risk of uterine rupture in women with more than 1 previous low transverse Caesarean section.^{8,78-84} All indicated success rates between 62% and 89%, and uterine rupture rates between 0% and 3.7%. In the largest study, Miller *et al.* demonstrated a VBAC success rate of 75.3% in 1827 women with 2 or more previous low transverse Caesarean sections with a uterine rupture rate of 1.7% vs. 0.6% in the ERCS group (OR, 3.06; 95% CI, 1.95–4.79).⁸ Unfortunately, the use of prostaglandins or oxytocin for induction or augmentation was not considered. Caughey *et al.* reported a uterine rupture rate of 3.7% versus 0.8% (RR, 4.8; 95% CI, 1.8–13.2) in a retrospective review of 134 women undergoing labour after 2 previous Caesarean sections after correction for prostaglandin, oxytocin, and epidural use.⁸⁴

RECOMMENDATION

12. **The available data suggest that a trial of labour in women with more than one previous Caesarean section is likely to be successful but is associated with a higher risk of uterine rupture. (II-2B)**

MULTIPLE PREGNANCY

Seven studies have examined a total of 233 women attempting VBAC in multiple pregnancy.⁸⁵⁻⁹¹ All support a trial of VBAC in multiple pregnancy as being safe and effective, with success rates of 69% to 84%, and without increased maternal or fetal morbidity or mortality.⁸⁵⁻⁹¹ In one study, uterine dehiscence was noted in 1 woman on manual exploration after successful vaginal delivery of both twins, and no treatment was required.⁸⁶ Each of these studies examined a small number of women, however; and greater numbers would be required to detect rare outcomes such as uterine rupture and maternal and perinatal mortality.

RECOMMENDATION

13. **Multiple gestation is not a contraindication to a TOL after Caesarean. (II-2B)**

BREECH PRESENTATION

A large multicentre trial by Hannah *et al.* demonstrated that a planned Caesarean birth is associated with better perinatal and neonatal outcomes in breech presentation at term.⁹² This recommendation has been adopted by the SOGC and would therefore preclude a planned TOL after Caesarean in women presenting with a singleton fetus in breech presentation at term.^{92,93} Vaginal delivery of premature fetuses and the second twin were not addressed in the study and therefore no recommendations can be made in this regard. It would seem appropriate to consider these cases individually. External cephalic version is not contraindicated in a woman with a previous Caesarean birth.^{94,95}

DIABETES MELLITUS

In a retrospective cohort study, Coleman *et al.* examined the issue of TOL after Caesarean in women with gestational diabetes mellitus (GDM).⁹⁶ Coleman examined 156 women with GDM and planned TOL after Caesarean and compared them to non-GDM women attempting TOL after Caesarean. They reported that the success rate for VBAC of 64.1% in women with GDM was lower than the 77.2% of women without GDM ($P < 0.001$).⁹⁶ Maternal and fetal morbidities were comparable.⁹⁶ A retrospective study of TOL after Caesarean in women with pregestational or gestational diabetes found similar results.⁹⁷ Based on these studies, diabetes mellitus should not be considered a contraindication to TOL after Caesarean.

RECOMMENDATION

14. **Diabetes mellitus is not a contraindication to TOL after Caesarean. (II-2B)**

MACROSOMIA

In a study examining the outcome of 365 women who underwent a TOL after Caesarean, giving birth to neonates weighing more than 4000 g, Zelop *et al.* demonstrated a success rate of

60%, with no increase in maternal or fetal morbidity and no increase in the risk of uterine rupture.⁹⁸ These data supported previously reported findings by Flamm (success rate 58%)⁹⁹ and Phelan (success rate 67%).¹⁰⁰ In 2003, Elkousy *et al.* reported an examination of 9960 women with a previous Caesarean section planning a trial of labour further stratified by neonatal birth weights and birth history (primarily whether they had had a previous vaginal delivery and if it occurred before or after their Caesarean section).¹⁰¹ His results indicate that the likelihood of successful VBAC decreases with increasing birth weight and is lowest in those who have never had a successful vaginal birth.¹⁰¹ Based on these results, suspected macrosomia is not a contraindication to a TOL after Caesarean, although it may be associated with a lower chance of success.

RECOMMENDATION

15. Suspected fetal macrosomia is not a contraindication to a TOL after Caesarean. (II-2B)

INTERDELIVERY INTERVAL

Four studies have examined the relationship between the inter-delivery interval and the rate of successful VBAC and uterine rupture.¹⁰²⁻¹⁰⁵ Esposito *et al.* examined 23 cases of uterine rupture and compared them to 127 controls.¹⁰² There was an increased risk of uterine rupture with a short interpregnancy interval (<6 months between pregnancies; <15 months between deliveries) compared to controls (17.4% vs. 4.7%, $P=0.05$).¹⁰² Shipp *et al.* reviewed 311 women who underwent a TOL after Caesarean less than 18 months after their Caesarean section and compared them to 2098 women who underwent a TOL after Caesarean after more than 18 months.¹⁰³ The shorter interval was associated with a 3-fold increase in the risk of uterine rupture (2.25% vs. 1.05%; OR, 3.0; 95% CI, 1.2–7.2).¹⁰³ Huang *et al.* reviewed 1185 women undergoing a TOL after Caesarean and noted no difference in the success of vaginal delivery in those with a shorter interval of <19 months (79% vs. 85.5%, $P=0.12$), but they did note a significant difference in successful VBAC in women who underwent medical induction compared to spontaneous labour (14.3% vs. 86.1%, $P<0.01$).¹⁰⁴ Their study noted no difference in the rate of uterine rupture.¹⁰⁴ In 2002, Bujold *et al.* reported an observational study of 1527 women undergoing a planned TOL after Caesarean at different intervals from the index Caesarean delivery.¹⁰⁵ The rates of uterine rupture were as follows: <12 months, 4.8%; 13 to 24 months, 2.7%; 25 to 36 months, 0.9%; and >36 months, 0.9%.¹⁰⁵ After adjustment for such confounders as number of layers in the uterine closure, induction, oxytocin, and epidural use, the odds ratio for uterine rupture in a woman <24 months from her last delivery was 2.65 (95% CI, 1.08–6.46).¹⁰⁵

RECOMMENDATION

16. Women delivering within 18 to 24 months of a

Caesarean section should be counselled about an increased risk of uterine rupture in labour. (II-2B)

POSTDATISM

Three studies have examined postdatism and TOL after Caesarean.¹⁰⁶⁻¹⁰⁸ In 2 of these studies, the rate of successful VBAC and uterine rupture in women who delivered at less than 40 weeks' gestation was compared to those who delivered at greater than 40 weeks.^{106,107} Success rates for VBAC after 40 weeks were reported from 65.6%¹⁰⁷ to 73.1%¹⁰⁶ and were comparable to success rates for women delivered before 40 weeks.^{106,107} Zelop *et al.* also compared the risk of uterine rupture in women who delivered before and after 40 weeks' gestation in spontaneous labour and induced labour.¹⁰⁸ They reported that the risk of uterine rupture in a TOL after Caesarean after 40 weeks was not significantly increased when compared to those who delivered before 40 weeks, whether in spontaneous labour (1.0% vs. 0.5%, $P=0.2$, adjusted OR, 2.1; 95% CI, 0.7–5.7) or after induction (2.6% vs. 2.1%, $P=0.7$, adjusted OR, 1.1; 95% CI, 0.4–3.4).¹⁰⁸

RECOMMENDATION

17. Postdatism is not a contraindication to a TOL after Caesarean. (II-2B)

ONE- VERSUS TWO-LAYER CLOSURE OF LOW TRANSVERSE CAESAREAN SECTION

In 1992, Hauth *et al.* published data comparing operative time, endometritis, transfusion, and placement of extra hemostatic sutures in women undergoing uterine closure in 1 layer compared to 2 layers.¹⁰⁹ The only significant difference was in operative time: 44 minutes with 1-layer compared to 48 minutes with 2-layer closure ($P<0.05$).¹⁰⁹ Similar findings were published by Ohel *et al.* in 1996.¹¹⁰ The trend shifted in many centres towards single-layer closure.

In 1997, Chapman *et al.* published a review of 145 women who underwent a TOL after Caesarean after being randomized to either 1-layer or 2-layer closure in the previous Caesarean section.¹¹¹ They reported no significant difference in the outcome of the next pregnancy.¹¹¹ In a 2002 review of 2142 women who underwent a TOL after Caesarean, Bujold *et al.* noted that a 1-layer interlocking closure was associated with an increased risk of uterine rupture when compared to a 2-layer closure (3.1% vs. 0.5%, $P<0.001$; OR, 3.95; 95% CI, 1.35–11.49).¹¹² Further study in this area is recommended.

UNKNOWN SCAR

All records available or obtainable describing the woman's previous Caesarean section should be reviewed. If unavailable, information about the circumstances of the Caesarean section will help determine the likelihood of a vertical uterine incision.^{113,114} The majority of unknown scars will be lower

transverse incisions (92%) and therefore at low risk for uterine rupture.¹¹⁵ If the history suggests a reasonable likelihood of a classical incision, it would be prudent to recommend a repeat Caesarean section, but in settings where the history indicates a high likelihood of lower transverse uterine incision and the woman wishes to proceed after counselling, a TOL after Caesarean is acceptable.¹¹⁵

RECOMMENDATION

18. Every effort should be made to obtain the previous Caesarean section operative report to determine the type of uterine incision used. In situations where the scar is unknown, information concerning the circumstances of the previous delivery is helpful in determining the likelihood of a low transverse incision. If the likelihood of a lower transverse incision is high, TOL after Caesarean can be offered. (II-2B)

OTHER FACTORS

Factors such as maternal obesity,¹¹⁶ the presence of postpartum fever after Caesarean section,¹¹⁷ type of suture material, müllerian duct anomalies,¹¹⁸ and maternal age¹¹⁹ and their relationship to the risk of uterine rupture have been examined in small studies, but definitive conclusions cannot yet be drawn.

CONCLUSION

Trial of labour after Caesarean section should be considered in women who have no contraindications after appropriate discussion. The efficacy and safety of a TOL after Caesarean in appropriately selected parturients in a hospital where immediate Caesarean section facilities are available is well supported. Support of the woman in labour, including close observation of her and her fetus for signs of complications, is recommended.

Augmentation of labour with oxytocin is safe. Induction of labour may be provided when the indication for induction is compelling and the risks have been fully discussed with the woman. The use of prostaglandin E₂ (dinoprostone) and prostaglandin E₁ (misoprostol) in women planning a TOL is not recommended. The use of a foley catheter for cervical ripening in situations where the cervix is unfavourable is associated with a lower chance of success but no increased risk of uterine rupture.

REFERENCES

- McMahon MJ, Luthier ER, Bowes WA, Olshan AF. Comparison of a trial of labor with an elective second Caesarean section. *N Engl J Med* 1996; 335:689–95.
- Davies GA, Hahn PM, McGrath MM. Vaginal birth after Caesarean section: physicians' perceptions and practice. *J Reprod Med* 1996;41: 515–20.
- Society of Obstetricians and Gynaecologists of Canada. Vaginal birth after previous Caesarean birth. Clinical Practice Guideline No. 68. Ottawa (ON): SOGC; December 1997.
- Biswass A. Management of previous Caesarean section. *Curr Opin Obstet Gynecol* 2003;15:123–9.
- Curtin SC, Kozak LJ, Gregory KD. U.S. Caesarean and VBAC rates stalled in the mid-1990s. *Birth* 2000;27:54–7.
- Health Canada. Canadian perinatal health report 2003. Ottawa (ON): Health Canada; 2003. p. 33.
- Weinstein D, Benshushan A, Ezra Y, Rojansky N. Vaginal birth after Caesarean section: current opinion. *Int J Gynecol Obstet* 1996;53:1–10.
- Miller DA, Diaz FG, Paul RH. Vaginal birth after Caesarean: a 10-year experience. *Obstet Gynecol* 1994;84:255–8.
- Schell JT. Once a Caesarean always a Caesarean? *NY Med J* 1923;637.
- Roberts LJ, Beardsworth SA, Trew G. Labour following Caesarean section: current practice in the United Kingdom. *Br J Obstet Gynaecol* 1994;101:153–5.
- Norman P, Kostovcik S, Lanning A. Elective repeat Caesarean sections: how many could be vaginal births? *Can Med Assoc J* 1993;149:431–5.
- Kline J, Arias F. Analysis of factors determining the selection of repeated Caesarean section or trial of labor in patients with histories of prior Caesarean delivery. *J Reprod Med* 1993;38:289–92.
- National Institutes of Health. Caesarean childbirth. Bethesda, MD: NIH; 1981. Publication No. 82-2067.
- Flamm BL, Lim OW, Jones C, Fallon D, Newman LA, Mantis JK. Vaginal birth after Caesarean section: results of a multicenter study. *Am J Obstet Gynecol* 1988;158:1079–84.
- Socol ML, Peaceman AM. Vaginal birth after Caesarean: an appraisal of fetal risk. *Obstet Gynecol* 1999;93:674–9.
- McMahon MJ. Vaginal birth after Caesarean. *Clin Obstet Gynecol* 1998; 41:369–81.
- Woolf SH, Battista RN, Angerson GM, Logan AG, Eel W. Canadian Task Force on the Periodic Health Exam. Ottawa (ON): Canada Communication Group; 1994. p. xxxvii.
- Quilligan EJ. Vaginal birth after Caesarean section: 270 degrees. *J Obstet Gynecol Res* 2001;27:169–73.
- Scott JR. Avoiding labor problems during vaginal birth after Caesarean delivery. *Clin Obstet Gynecol* 1997;40:533–41.
- Rageth JC, Juzi C, Grossenbacher H. Delivery after previous Caesarean: a risk evaluation. *Obstet Gynecol* 1999;93:332–7.
- Lovell R. Vaginal delivery after Caesarean section: factors influencing success rates. *Aust N Z J Obstet Gynaecol* 1996;36:4–8.
- Weinstein D, Benshushan A, Tanos V, Zilberstein R, Rojansky N. Predictive score for vaginal birth after Caesarean section. *Am J Obstet Gynecol* 1996;174:192–8.
- Flamm BL, Geiger AM. Vaginal birth after Caesarean delivery: an admission scoring system. *Obstet Gynecol* 1997;90:907–10.
- Bujold E, Gauthier R. Should we allow a trial of labor after a previous Caesarean for dystocia in the second stage of labor? *Obstet Gynecol* 2001;98:652–5.
- Hoskins IA, Gomez JL. Correlation between maximum cervical dilatation at Caesarean delivery and subsequent vaginal birth after Caesarean delivery. *Obstet Gynecol* 1997;89:591–3.
- Shipp TD, Zelop CM, Repke JT, Cohen A, Caughey AB, Lieberman E. Labor after previous Caesarean: influence of prior indication and parity. *Obstet Gynecol* 2000;95:913–6.
- Hibbard JU, Ismail MA, Wang Y, Te C, Karrison T, Ismail MA. Failed vaginal birth after Caesarean section: how risky is it? *Am J Obstet Gynecol* 2001;184:1365–73.
- Rosen MG, Dickinson JC, Westhoff CL. Vaginal birth after Caesarean section: a meta-analysis of morbidity and mortality. *Obstet Gynecol* 1991;77:465–70.
- Hemminki E, Merilainen J. Long-term effects of Caesarean sections: ectopic pregnancies and placental problems. *Am J Obstet Gynecol* 1996;174:1569–74.
- Gilliam M, Rosenberg D, Davis D. The likelihood of placenta previa with greater number of Caesarean deliveries and higher parity. *Obstet Gynecol* 2002;99:976–80.
- Miller DA, Chollet JA, Goodwin TM. Clinical risk factors for placenta previa-placenta accreta. *Am J Obstet Gynecol* 1997;177:210–4.
- Mozurkewich EL, Hutton EK. Elective repeat Caesarean delivery versus

- trial of labor: a meta-analysis of the literature from 1989 to 1999. *Am J Obstet Gynecol* 2000;183:1187-97.
33. Hook B, Kiwi R, Amini SB, Fanaroff A, Hack M. Neonatal morbidity after elective repeat Cesarean section and trial of labor. *Pediatrics* 1997;100:348-53.
 34. American College of Obstetricians and Gynecologists. Vaginal birth after previous Cesarean delivery. ACOG Practice Bulletin No. 5. Washington (DC): ACOG; 1999.
 35. Wing DA, Paul RH. Vaginal birth after Cesarean section: selection and management. *Clin Obstet Gynecol* 1999;42:836-48.
 36. Khan KS, Risvi A. The partograph in the management of labor following Cesarean section. *Int J Gynecol Obstet* 1995;50:151-7.
 37. Guleria K, Dhall GI, Dhall K. Pattern of cervical dilatation in previous lower segment Cesarean section patients. *J Indian Med Assoc* 1997;95:131-4.
 38. Rowbottom SJ, Critchley LAH, Gin T. Uterine rupture and epidural analgesia during trial of labor. *Anaesthesia* 1997;52:483-8.
 39. Society of Obstetricians and Gynaecologists of Canada. Fetal health surveillance in labour. *J Obstet Gynaecol Can* 2002;112:250-62.
 40. Phelan JP, Korst LM, Settles DK. Uterine activity patterns in uterine rupture: a case-control study. *Obstet Gynecol* 1998;92:394-7.
 41. Kaplan B, Royburt M, Peled Y, Hirsch M, Ovadia Y, Neri A. Routine revision of uterine scar after prior Cesarean section. *Acta Obstet Gynecol Scand* 1994;73:473-5.
 42. Bucklin BA. Vaginal birth after Cesarean delivery. *Anesthesiology* 2003;99:1444-8.
 43. Hill DA, Chez RA, Quinlan J, Fuentes A, LaCombe J. Uterine rupture and dehiscence associated with intravaginal misoprostol cervical ripening. *J Reprod Med* 2000;45:823-6.
 44. Yap OVS, Kim ES, Laros RK. Maternal and neonatal outcomes after uterine rupture in labor. *Am J Obstet Gynecol* 2001;184:1576-81.
 45. Appleton B, Targett C, Rasmussen M, Readman E, Sale F, Permezel M. Vaginal birth after Caesarean section: an Australian multicentre study. *Aust N Z J Obstet Gynaecol* 2000;40:87-91.
 46. Shipp TA, Zelop CM, Repke JT, Cohen A, Caughey AB, Lieberman E. Intrapartum uterine rupture and dehiscence in patients with prior lower uterine segment vertical and transverse incisions. *Obstet Gynecol* 1993;94:735-40.
 47. Martin JN, Perry KG, Roberts WE, Meydrech EF. The case for trial of labor in the patient with a prior low-segment vertical Cesarean incision. *Am J Obstet Gynecol* 1997;177:144-8.
 48. Adair CD, Sanchez-Ramos L, Whitaker D, McDyer DC, Farah L, Briones D. Trial of labor in patients with a previous lower uterine vertical Cesarean section. *Am J Obstet Gynecol* 1996;174:966-70.
 49. Stovall TG, Shaver DC, Solomon SK, Anderson GD. Trial of labor in previous Cesarean section patients, excluding classical Cesarean sections. *Obstet Gynecol* 1987;70:713-7.
 50. Shimonovitz S, Botosneano A, Hochner-Celnikier D. Successful first vaginal birth after Cesarean section: a predictor of reduced risk for uterine rupture in subsequent deliveries. *Indian Med Assoc J* 2000;2:526-8.
 51. Kieser KE, Baskett TF. A 10-year population-based study of uterine rupture. *Am J Obstet Gynecol* 2002;100:749-53.
 52. Bujold E, Gauthier RJ. Neonatal morbidity associated with uterine rupture: what are the risk factors? *Am J Obstet Gynecol* 2002;186:311-4.
 53. Smith GCS, Pell JP, Cameron AD, Dobbie R. Risk of perinatal death associated with labor after previous Cesarean delivery in uncomplicated term pregnancies. *J Am Med Assoc* 2002;287:2684-90.
 54. Chauhan SP, Martin JN, Henrichs CE, Morrison JC, Magann EF. Maternal and perinatal complications with uterine rupture in 142 075 patients who attempted vaginal birth after Cesarean delivery: a review of the literature. *Am J Obstet Gynecol* 2003;189:408-17.
 55. Hamilton EF, Bujold E, McNamara H, Gauthier R, Platt RV. Dystocia among women with symptomatic uterine rupture. *Am J Obstet Gynecol* 2001;184:620-4.
 56. Rozenberg P, Goffinet F, Philippe HJ, Nisand I. Ultrasonographic measurement of lower uterine segment to assess the risk of defects of scarred uterus. *Lancet* 1996;347:281-4.
 57. Rozenberg P, Goffinet F, Philippe HJ, Nisand I. Thickness of the lower uterine segment: its influence in the management of patients with previous Cesarean sections. *Eur J Obstet Gynecol Reprod Biol* 1999;87:39-45.
 58. Flamm BL, Goings JR, Fielbirth N, Fischermann E, Jones C, Hewson SA. Oxytocin during labor after previous Cesarean section: results of a multicenter study. *Obstet Gynecol* 1987;70:709-12.
 59. Zelop CM, Shipp TA, Repke JT, Cohen A, Lieberman E. Uterine rupture during induced or augmented labor in gravid women with one prior Cesarean delivery. *Am J Obstet Gynecol* 1999;181:882-6.
 60. Goetzl L, Shipp TA, Cohen A, Zelop CM, Repke JT, Lieberman E. Oxytocin dose and the risk of uterine rupture in trial of labor after Cesarean. *Obstet Gynecol* 2001;97:381-4.
 61. Ravasia DJ, Wood SL, Pollard JK. Uterine rupture during induced trial of labor among women with previous Cesarean delivery. *Am J Obstet Gynecol* 2000;183:1176-9.
 62. Sanchez-Ramos L, Gaudier FL, Kaunitz AM. Cervical ripening and labor induction after previous Cesarean delivery. *Clin Obstet Gynecol* 2000;43:513-23.
 63. Delaney T, Young DC. Spontaneous versus induced labor after a previous Cesarean delivery. *Obstet Gynecol* 2003;102:39-44.
 64. Stone JL, Lockwood CJ, Berkowitz G, Alvarez M, Lapinski R, Valcamonica A, et al. Use of cervical prostaglandin E₂ gel in patients with previous Cesarean section. *Am J Perinatol* 1994;11:309-12.
 65. Flamm BL, Anton D, Goings JR, Newman J. Prostaglandin E₂ for cervical ripening: a multicenter study of patients with prior Cesarean delivery. *Am J Perinatol* 1997;14:157-60.
 66. MacKenzie JZ, Bradley S, Embrey MP. Vaginal prostaglandins and labour induction for patients previously delivered by Caesarean section. *Br J Obstet Gynaecol* 1984;91:7-10.
 67. Lyndon-Rochelle M, Holt VL, Easterling TR, Martin DP. Risk of uterine rupture during labor among women with a prior Cesarean delivery. *N Engl J Med* 2001;345:3-8.
 68. Katz VL, Farmer RM, Dean CA, Carpenter ME. Use of misoprostol for cervical ripening. *Southern Med J* 2000;93:881-4.
 69. Sciscione AC, Nguyen L, Manley JS, Schlossman PA, Colmorgen GHC. Uterine rupture during preinduction cervical ripening with misoprostol in a patient with a previous Caesarean delivery. *Aust N Z J Obstet Gynaecol* 1998;38:96-7.
 70. Choy-Hee L, Raynor BD. Misoprostol induction of labor among women with a history of Cesarean delivery. *Am J Obstet Gynecol* 2001;184:1115-7.
 71. Cunha M, Bugalho A, Bique C, Berstrom S. Induction of labor by vaginal misoprostol in patients with previous Cesarean delivery. *Acta Obstet Gynecol Scand* 1999;78:653-4.
 72. Plaut MM, Schwartz ML, Lubarsky SL. Uterine rupture associated with the use of misoprostol in the gravid patient with a previous Cesarean section. *Am J Obstet Gynecol* 1999;180:1535-42.
 73. Wing DA, Lovett K, Paul RH. Disruption of prior uterine incision following misoprostol for labor induction in women with previous Cesarean delivery. *Obstet Gynecol* 1998;91:828-30.
 74. Blanchette HA, Nayak S, Erasmus S. Comparison of the safety and efficacy of intravaginal misoprostol (prostaglandin E₂) with that of dinoprostone (prostaglandin E₂) for cervical ripening and induction of labor in a community hospital. *Am J Obstet Gynecol* 1999;180:1551-9.
 75. American College of Obstetricians and Gynecologists. Induction of labor for vaginal birth after Cesarean delivery. *Obstet Gynecol* 2002;99:679-80.
 76. Ben-Aroya Z, Hallak M, Segal D, Friger M, Katz M, Mazor M. Ripening of the uterine cervix in a post-Cesarean parturient: prostaglandin E₂ versus foley catheter. *J Matern Fetal Neonatal Med* 2002;12:42-5.
 77. Bujold E, Blackwell SC, Gauthier RJ. Cervical ripening with transcervical foley catheter and the risk of uterine rupture. *Obstet Gynecol* 2004;103:18-23.
 78. Asakura H, Myers SA. More than one previous Cesarean delivery: a 5-year experience with 435 patients. *Obstet Gynecol* 1995;85:924-9.
 79. Chattopadhyay SK, Sherbeeni MM, Anokute CC. Planned vaginal delivery after two previous Caesarean sections. *Br J Obstet Gynaecol* 1994;101:498-500.

80. Bretelle F, Cravello L, Shojai R, Roger V, D'Ercole C, Blanc B. Vaginal birth following two previous Cesarean sections. *Eur J Obstet Gynecol Reprod Biol* 2001;94:23–6.
81. Phelan JP, Ahn MO, Diaz F, Brar HS, Rodriguez MH. Twice a Cesarean, always a Cesarean? *Obstet Gynecol* 1989;73:161–5.
82. Porreco RP, Meier PR. Trial of labor in patient with multiple previous Cesarean sections. *J Reprod Med* 1983;28:770–2.
83. Farmakides G, Duvioler R, Schulman H, Schneider E, Biordi J. Vaginal birth after two or more previous Cesarean sections. *Am J Obstet Gynecol* 1987;156:565–6.
84. Caughey AB, Shipp TA, Repke JT, Zelop CM, Cohen A, Lieberman E. Rate of uterine rupture during a trial of labor in women with one or two prior Cesarean deliveries. *Am J Obstet Gynecol* 1999;181:872–6.
85. Miller DA, Mullin P, Hou D, Paul RH. Vaginal birth after Cesarean section in twin gestation. *Am J Obstet Gynecol* 1996;175:194–8.
86. Strong TH, Phelan JP, Ahn MO, Sarno AP. Vaginal birth after Cesarean section in the twin gestation. *Am J Obstet Gynecol* 1989;161:29–32.
87. Odeh M, Tarazova L, Wolfson M, Oettinger M. Evidence that women with a history of Cesarean section can deliver twins safely. *Acta Obstet Gynecol Scand* 1997;76:663–6.
88. Myles T. Vaginal birth of twins after previous Cesarean section. *J Maternal Fetal Med* 2001;10:171–4.
89. Wax JR, Philput C, Mather J, Steinfeld JD, Ingardia CJ. Twin vaginal birth after Cesarean. *Conn Med* 2000;64:205–8.
90. Sansregret A, Bujold E, Gauthier RJ. Twin delivery after a previous Cesarean: a twelve-year experience. *J Obstet Gynaecol Can* 2003;25:294–8.
91. Delaney T, Young DC. Trial of labour compared to elective Cesarean in twin gestations with a previous Cesarean delivery. *J Obstet Gynaecol Can* 2003;25:289–92.
92. Hannah ME, Hannah WJ, Hewson SA. Planned Cesarean section versus planned vaginal birth for breech presentation at term: a randomised multicenter trial. *Lancet* 2000;356:1375–83.
93. Society of Obstetricians and Gynaecologists of Canada. SOGC statement on vaginal breech. Ottawa (ON): SOGC; 1999.
94. de Meeus JB, Elia F, Magnin G. External cephalic version after previous Cesarean section: a series of 38 cases. *Eur J Obstet Gynecol Reprod Biol* 1998;81:65–8.
95. Flamm BL, Fried MW, Lonky NM. External cephalic version after previous Cesarean section. *Am J Obstet Gynecol* 1991;165:370–2.
96. Coleman TL, Randall H, Graves W, Lindsay M. Vaginal birth after Cesarean among women with gestational diabetes. *Am J Obstet Gynecol* 2001;184:1104–7.
97. Blackwell SC, Hassan SS, Wolfe HM, Michaelson J, Berry SM, Sorokin Y. Vaginal birth after Cesarean in the diabetic gravida. *J Reprod Med* 2000;45:987–90.
98. Zelop CM, Shipp TA, Repke JT, Cohen A, Lieberman E. Outcomes of trial of labor following previous Cesarean delivery among women with fetuses weighing >4000 g. *Am J Obstet Gynecol* 2001;185:903–5.
99. Flamm BL, Goings JR. Vaginal birth after Cesarean section: is suspected fetal macrosomia a contraindication? *Obstet Gynecol* 1989;74:694–7.
100. Phelan JP, Eglinton GS, Horenstein JM. Previous Cesarean birth: trial of labor in women with macrosomic infants. *J Reprod Med* 1984;29:36–40.
101. Elkousy MA, Sammel M, Stevens E, Peipert JF, Macones G. The effect of birth weight on vaginal birth after Cesarean delivery success rates. *Am J Obstet Gynecol* 2003;188:824–30.
102. Esposito MA, Menihan CA, Malee MP. Association of interpregnancy interval with uterine scar failure in labor: a case-control study. *Am J Obstet Gynecol* 2000;183:1180–3.
103. Shipp TA, Zelop CM, Repke JT, Cohen A, Lieberman E. Interdelivery interval and risk of symptomatic uterine rupture. *Obstet Gynecol* 2001;97:175–7.
104. Huang WH, Nakashima DK, Rumney PJ, Keegan KA, Chan K. Interdelivery interval and the success of vaginal birth after Cesarean delivery. *Obstet Gynecol* 2002;99:41–4.
105. Bujold E, Mehta SH, Bujold C, Gauthier R. Interdelivery interval and uterine rupture. *Am J Obstet Gynecol* 2002;187:1199–202.
106. Yeh S, Huang X, Phelan JP. Post-term pregnancy after previous Cesarean section. *J Reprod Med* 1984;29:41–4.
107. Callahan C, Chescheir N, Steiner BD. Safety and efficacy of attempted vaginal birth after Cesarean beyond the estimated date of delivery. *J Reprod Med* 1999;44:606–10.
108. Zelop CM, Shipp TA, Cohen A, Repke JT, Lieberman E. Trial of labor after 40 weeks' gestation in women with prior Cesarean. *Obstet Gynecol* 2001;97:391–3.
109. Hautz JC, Owen J, Davis RO. Transverse uterine incision closure: one versus two layers. *Am J Obstet Gynecol* 1992;167:1108–11.
110. Ohel G, Younis JS, Lang N, Levit A. Double-layer closure of uterine incision with visceral and parietal peritoneal closure: are they obligatory? *J Matern Fetal Med* 1996;5:366–9.
111. Chapman SJ, Owen J, Hautz JC. One- versus two-layer closure of a low transverse Cesarean: the next pregnancy. *Obstet Gynecol* 1997;89:16–8.
112. Bujold E, Bujold C, Hamilton EF, Harel F, Gauthier R. The impact of a single-layer or double-layer closure on uterine rupture. *Am J Obstet Gynecol* 2002;186:1326–30.
113. Grubb DK, Kjos SL, Paul RH. Latent labor with an unknown uterine scar. *Obstet Gynecol* 1996;88:351–5.
114. Lau TK, Chan F. Unknown uterine scars, unknown risks. *Aust N Z J Obstet Gynaecol* 1994;34:216–7.
115. Beall M, Eglinton GS, Clark SL, Phelan JP. Vaginal delivery after Cesarean section in women with unknown types of uterine scar. *J Reprod Med* 1984;29(1):32–5.
116. Chauhan SP, Magann EF, Carroll CS, Barrilleaux PS, Scardo JA, Martin JN. Mode of delivery for the morbidly obese with prior Cesarean delivery: vaginal versus repeat Cesarean section. *Am J Obstet Gynecol* 2001;185:349–54.
117. Shipp TA, Zelop CM, Cohen A, Repke JT, Lieberman E. Post-Cesarean delivery fever and uterine rupture in a subsequent trial of labor. *Am J Obstet Gynecol* 2003;101:136–9.
118. Ravasia DJ, Brain PH, Pollard JK. Incidence of uterine rupture among women with müllerian duct anomalies who attempt vaginal birth after Cesarean delivery. *Am J Obstet Gynecol* 1999;181:877–81.
119. Shipp TA, Zelop CM, Repke JT, Cohen A, Caughey AB, Lieberman E. Maternal age as a predictor of symptomatic uterine rupture during a trial of labor after previous Cesarean delivery. *Am J Obstet Gynecol* 2001;184:S186.